

## Effects of dihydrocubebin, a lignan isolated from Indonesian plant *Piper cubeba*, on the histamine release from rat mast cells

Agung Endro Nugroho<sup>1,2,\*</sup>, Wahyuono<sup>3</sup>, Subagus Wahyuono<sup>3</sup> and Kazutaka Maeyama<sup>1</sup>

<sup>1</sup>Department of Pharmacology, Informational Biomedicine, Ehime University Graduate School of Medicine, Shitsugawa, Toon-shi, Ehime 791-0295, Japan; <sup>2</sup>Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Gadjah Mada University, Jogjakarta 55281, Indonesia; <sup>3</sup>Department of Pharmaceutical Biology, Faculty of Pharmacy, Gadjah Mada University, Jogjakarta, 55281, Indonesia

Received for publication February 25 2009; accepted September 7 2010

### SUMMARY

The fruits of *Piper cubeba* L. are used traditionally to treat respiratory disorders in Indonesia. In order to determine the compounds responsible for this activity, the fruits were extracted with n-hexane followed by ethanol to give n-hexane and ethanol extracts. Based on tracheospasmolytic assay on these two extracts, the n-hexane extract was more active to inhibit trachea contraction than that of ethanol extract. Upon bioassay guided isolation of the n-hexane extract, a tracheospasmolytic active compound was isolated and identified as dihydrocubebin [(3,4),(3',4')-bis-methylenedioxy-9,9-dihydroxylignan] (**1**). Compound **1** was tested further for its ability to inhibit histamine released from mast cells, using rat basophilic leukemia (RBL-2H3) cell line and rat peritoneal mast cells (RPMCs) as models; and DNP<sub>24</sub>-BSA, thapsigargin, ionomycin, compound 48/80 and PMA were used as inducers for histamine released from mast cell. The test result showed that **1** inhibited histamine release from RBL-2H3 cells induced by DNP<sub>24</sub>-BSA, thapsigargin and ionomycin. In addition, **1** suppressed histamine release from RPMC induced by either thapsigargin or ionomycin. However, **1** did not inhibit histamine release from RPMC induced by either compound 48/80 or combination PMA-sub optimum dose of ionomycin. Therefore, it was concluded that the inhibitory effects of **1** on the histamine released from mast cells may involve mechanisms related to intracellular Ca<sup>2+</sup> signaling events or downstream processes of intracellular Ca<sup>2+</sup> signaling in mast cells.

**Key words:** *Piper cubeba* L.; Dihydrocubebin; Histamine; Mast cells

### INTRODUCTION

Indonesia has been reported for its first biodiversity of the world. *Piper cubeba* L. is commonly present in a traditional preparation (*jamu*) intended to cure respiratory disorder. Initial study indicated that n-

hexane extract of *Piper cubeba* inhibited contraction of isolated guinea pig trachea induced by methacholine (Wahyuono *et al.*, 1999). Many plants of genus *Piper* have been studied for their anti-allergy effects such as of *Piper longum* using rat-lung perfusion as a model (Nayampali and Satoskar, 1980). Ethanolic extract of *Piper betle* leaves decreased histamine and GM-CSF produced by an IgE-mediated hypersensitive reaction, and inhibited eotaxin and IL-8 secretion as well in a TNF- $\alpha$  and IL-4-induced allergic reaction

\*Correspondence: Agung Endro Nugroho, Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Gadjah Mada University, Jogjakarta 55281, Indonesia. Tel: +0274543120; Fax: +0274543120; E-mail: agungendronugroho@yahoo.com, nugroho\_ae@ugm.ac.id

(Wirotasangthong *et al.*, 2007). While methanolic extract of *Piper nigrum* leaves inhibited histamine release from rat peritoneal mast cell, a powerful inhibition on DNFB-induced cutaneous reaction at 1 h and 24 h after DNFB challenge in mice, which passively sensitized with anti-DNP IgE antibody upon oral administration of this extract (Hirata *et al.*, 2008).

Asthma is a lung disease characterized by airways obstruction, airways inflammation, and increased airway responsiveness. Allergic reaction has a main role on development of airway obstruction and pathogenesis of asthma (Busse and Reed, 1993; Holt *et al.*, 1999). Mast cell is a crucial effector cell in the pathogenesis of asthma, especially asthma with an allergic basis (Brown *et al.*, 2008). Mast cells are strategically located in pulmonary system to participate in lung allergic reactions by release rapidly mediators such as histamine and other lipid mediators (Busse and Reed, 1993). A therapeutic strategy for treating allergic airway diseases is to inhibit mast cells degranulation and to inhibit the release of mast cells mediators such as histamine.

At present we report the effect of dihydrocubebin [(3,4),(3',4')-bis-methylenedioxy-9,9'-dihydroxylignan] (1), a lignan isolated from *Piper cubeba* fructus (Prabhu and Mulchandani, 1985; Wahyono, 2005) to inhibit histamine released from mast cells. In the study, rat basophilic leukemia (RBL-2H3) cells and rat peritoneal mast cells (RPMCs) were used; and DNP<sub>24</sub>-BSA, thapsigargin, ionomycin, compound 48/80 and PMA were used as inducers for histamine release from mast cells.

## MATERIALS AND METHODS

### Materials

Dihydrocubebin (1) was isolated from *Piper cubeba* L.f fruits by Dr. Wahyono from Gadjah Mada University (GMU), Indonesia. The histamine release inducers used in the study were ionomycin, thapsigargin, compound 48/80, and phorbol myristate

acetate (PMA) from Sigma Chemical Co. (St.Louis, MO, USA). Dinitro-phenylated bovine serum albumin (DNP<sub>24</sub>-BSA) as an antigen and monoclonal IgE against DNP<sub>24</sub>-BSA purified from supernatant in IgE producing hybridoma, were produced in our laboratory. Eagle's minimum essential medium (MEM) and antibiotics (combination of penicillin G sodium and streptomycin sulfate) were purchased from Gibco (Grand Island, NY, USA). Fetal calf serum was obtained from JRH Biosciences (Kansas, USA). Piperazine-1,4-bis(2-ethanesulfonic acid) (PIPES) was purchased from Dojindo (Kumamoto, Japan), and o-phthalaldehyde was from Wako Pure Chemical Industries (Osaka, Japan).

### Culture of RBL-2H3 cells

The RBL-2H3 cells were cultured in MEM containing 15% fetal calf serum and antibiotics (penicillin and streptomycin) in a flask in a humidified atmosphere (5% CO<sub>2</sub>) at 37 °C as described by Barsumian *et al.* (1981). For the assay of histamine release, cells were seeded into 24-well culture plates at a density of  $5 \times 10^5$  cells/0.4 ml per each well. The cells were incubated overnight at 37 °C. For DNP<sub>24</sub>-BSA experiments, the cells were sensitized with 0.5 µg/ml of monoclonal IgE. On the second day, the medium was removed, and the cells were washed twice with 500 µl of PIPES buffer (119 mM NaCl, 5 mM KCl, 25 mM PIPES, 5.6 mM glucose, 0.4 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub>, 40 mM NaOH, and 0.1% bovine serum albumin, pH 7.2), and pre-incubated for 10 min at 37 °C after addition of 180 µl PIPES buffer either without (as a negative control) or with the drug. After 10 min pre-incubation, 20 µl of stimulant (200 ng/ml DNP<sub>24</sub>-BSA, 5 µM thapsigargin, or 10 µM ionomycin) were added to each well and the plate was incubated at 37 °C for 30 min.

### Isolation of RPMCs

Male Wistar rats weighing between 250 - 300 g (3 - 4 months) were used. The animal experiments were conducted according to the guidelines of the Animal Care Committee of the Ehime University, and all

experimental protocols had been approved by this Committee. RPMCs were isolated by injection of 25 ml phosphate buffered saline (PBS) pH 7.4 containing 5 IU/ml heparin and 0.1% BSA into the peritoneal cavity and the abdomen was massaged for about 120 s. Afterwards, the peritoneal cavity was opened carefully, and the fluid containing mast cells were collected. The collected mast cells were centrifuged at 1,000 rpm for 5 min at room temperature and then re-suspended in 2 ml PBS buffer containing 0.1% BSA. Peritoneal mast cells were separated from the other components (macrophages and lymphocytes) by layering on 4 ml of 38% BSA, and centrifuging at 2,000 rpm for 20 min at 4 °C. After the upper layer containing other components was aspirated and discarded, the remaining cell pellet was washed with 6 ml PBS buffer and re-suspended in 1 ml of PIPES buffer. Mast cell preparations were about 95% pure as assessed by toluidine blue staining.

For the assay of histamine release, 120 µl of RPMCs suspension ( $2 \times 10^4$  cells/ml) was pre-incubated for 10 min at 37 °C after addition of 60 µl PIPES buffer either without (as a negative control) or with drugs at a range of concentrations (0.1 – 100 µM). After 10 min pre-incubation, 20 µl of stimulant (100 µg/ml compound 48/80, 5 µM thapsigargin, 10 µM ionomycin, or a combination of 100 nM PMA and 1 µM ionomycin) was added to each well and the plates were incubated at 37 °C for 30 min.

#### Assay of histamine release

Histamine released in the medium was measured by HPLC-fluorometry as described previously (Yamatodani *et al.*, 1985). After 30 min incubation, the plates were centrifuged at 3,000 rpm for 5 min and 50 µl of the supernatant was mixed with 250 µl of 3% perchloric acid containing 5 mM Na<sub>2</sub>-EDTA. After addition of 30 µl of 2 M KOH/1 M KH<sub>2</sub>PO<sub>4</sub> and centrifugation at 10,000 × g for 15 min at 4 °C, 50 µl of the supernatant was injected directly onto a column packed with TSKgel SP-2SW cation exchanger (Tosoh, Tokyo). For measuring the total histamine




Fig. 1. Dihydrocubebine (**1**).

content in cells, 350 µl of PIPES buffer was added to 6 wells and the cells were then sonicated. Fifty microlitres cell homogenate was used for the histamine assay described above. Histamine was eluted with 0.25 M potassium phosphate at a flow rate of 0.6 ml/min, and post-labeled with *o*-phthalaldehyde under alkaline conditions and detected using a F1080 Fluorometer (Hitachi, Tokyo) at excitation and emission wavelengths of 360 and 450 nm, respectively. The values were expressed as a percentage of net histamine release.


#### Statistical analysis

All data were expressed as mean ± SEM. One-way analysis of variance (ANOVA) followed by the least significant difference (LSD) test was used for statistical analysis. *P*-values less than 0.05 were considered significant.


## RESULTS

#### Effects on histamine release from RBL-2H3 cells

DNP<sub>24</sub>-BSA was used as an antigen to stimulate histamine release from IgE-sensitized RBL-2H3 cells. DNP<sub>24</sub>-BSA (20 ng/ml) stimulated histamine release from RBL-2H3 cells by  $34.56 \pm 1.05\%$  ( $n = 3$ ) (Fig. 2). Pre-incubation of RBL-2H3 cells with **1** for 30 minutes significantly inhibited the histamine and β-hexoaminidase enzyme release induced by DNP<sub>24</sub>-BSA in a dose-dependent manner. At the highest dose (100 µM), this compound inhibited the release of histamine from RBL-2H3 cells by  $52.05 \pm 3.85\%$  ( $n = 3$ ) (Table 1). IC<sub>50</sub> value of its



**Fig. 2.** Effect of dihydrocubebin on histamine released from RBL-2H3 cells in the presence of DNP-BSA 20 ng/ml. Data represent mean  $\pm$  SEM, and are three independent experiments. \* $P < 0.05$  compared to the negative control value.



**Fig. 3.** Effect of dihydrocubebin on histamine released from RBL-2H3 cells in the presence of either thapsigargin 0.5  $\mu$ M (a); or ionomycin 1  $\mu$ M (b). Data represent mean  $\pm$  SEM, and are three independent experiments. \* $P < 0.05$ , \*\* $P < 0.1$  compared to the negative control value.

inhibitory effect was used as the parameter of potency. The  $IC_{50}$  of the inhibitory effects of this compound on histamine induced by DNP<sub>24</sub>-BSA

**Table 1.** The inhibitory effects of dihydrocubebin on the histamine released from RBL-2H3 cells and its  $IC_{50}$  values

No.	Conc. ( $\mu$ M)	histamine release inhibition (%)		
		DNP <sub>24</sub> -BSA	thapsigargin	ionomycin
1.	1	0.78 $\pm$ 9.71	6.84 $\pm$ 10.67	2.51 $\pm$ 8.19
2.	3	6.05 $\pm$ 6.38	19.38 $\pm$ 8.91	5.88 $\pm$ 9.68
3.	10	26.00 $\pm$ 0.49	30.25 $\pm$ 1.86	18.69 $\pm$ 3.51
4.	30	30.51 $\pm$ 5.93	37.64 $\pm$ 2.58	29.27 $\pm$ 2.22
5.	100	52.05 $\pm$ 3.85	59.75 $\pm$ 1.48	61.35 $\pm$ 8.88
$IC_{50}$ value ( $\mu$ M)		93.34	69.13	75.23

Concentration of DNP<sub>24</sub>-BSA, thapsigargin and ionomycin were 20 ng/ml, 0.5  $\mu$ M and 1  $\mu$ M, respectively. Data represent mean  $\pm$  SEM of three independent experiments.


were 93.34  $\mu$ M.

Thapsigargin (0.5  $\mu$ M) and ionomycin (1  $\mu$ M), which act on calcium ion influx and intracellular calcium pathways, stimulated histamine release by 36.44  $\pm$  1.54% ( $n = 3$ ); and 30.22  $\pm$  1.99% ( $n = 3$ ), respectively (Fig. 3). Dihydrocubebin (1) succeeded to inhibit the histamine release induced by both of which in a dose-dependent manner. At the dose of 100  $\mu$ M, the compound strongly inhibited the release of histamine from RBL-2H3 cells by 59.75  $\pm$  1.48% ( $n = 3$ ); and 61.35  $\pm$  8.88% ( $n = 3$ ), respectively (Table 1). The  $IC_{50}$  of the inhibitory effects of 1 on histamine release induced by both inducers were 69.13 and 75.23  $\mu$ M respectively.

In the absence of histamine secretagogue, possibility of histamine release from RBL-2H3 cells by 1 was observed. The effect was considered significant if the compound caused spontaneous histamine release of more than 10% (Ikawati *et al.*, 2001). Dihydrocubebin at 100  $\mu$ M induced the spontaneous histamine release on RBL-2H3 cells by 1.08  $\pm$  0.97% ( $n = 3$ ).

#### Effects on histamine release from RPMCs

Compound 48/80, PMA, thapsigargin, and ionomycin were used for stimulating the histamine release rat peritoneal mast cells (RPMCs). Compounds 48/80, which acts on G proteins in mast cells, stimulated histamine release from RPMCs by 73.26  $\pm$  1.41% ( $n$




**Fig. 4.** Effect of dihydrocubebin on histamine released from rat peritoneal mast cells in the presence of either compound 48/80 10 µg/ml (a); or phorbol myristate acetate 10 nM-ionomycin 0.1 µM (b). Data represent mean  $\pm$  SEM, and are three independent experiments.

= 3). Dihydrocubebin at 0.1-100 µM did not significantly inhibit compound 48/80-induced histamine release.

To assess the effect of **1** on protein kinase C-mediated histamine release, combination of PMA, a modulator of PKC, and low-dose ionomycin were used. RPMCs release  $90.93 \pm 0.66\%$  ( $n = 3$ ) of their total histamine content in response to combination 10 nM PMA and 0.1 µM ionomycin (Fig. 4). In this experiment, **1** (0.1 - 100 µM) did not significantly influence the histamine release.

In RPMCs, thapsigargin (0.5 µM) and ionomycin (1 µM) induced histamine release by  $54.02 \pm 0.97\%$  ( $n = 3$ ); and  $79.52 \pm 0.60\%$  ( $n = 3$ ), respectively (Fig. 5). Dihydrocubebin dose-dependently inhibited both thapsigargin- and ionomycin-induced histamine released from RPMC. At the dose of 100 µM, the compound potently inhibited the release of histamine release from RBL-2H3 cells by  $60.65 \pm 1.21\%$  ( $n = 3$ ); and  $23.23 \pm 7.19\%$  ( $n = 3$ ), respectively (Table 2). The



**Fig. 5.** Effect of dihydrocubebin on histamine released from rat peritoneal mast cells in the presence of either thapsigargin 0.5 µM (a); or ionomycin 1 µM (b). Data represent mean  $\pm$  SEM, and are three independent experiments. \* $P < 0.05$  compared to the negative control value.

**Table 2.** The inhibitory effects of dihydrocubebin on the histamine release from RBL-2H3 cells

No.	Conc. (mM)	histamine release inhibition (%)	
		thapsigargin	ionomycin
1.	1	$0.85 \pm 2.24$	$0.47 \pm 1.61$
2.	3	$1.89 \pm 4.26$	$0.40 \pm 1.22$
3.	10	$5.13 \pm 4.34$	$2.22 \pm 4.12$
4.	30	$17.52 \pm 0.61$	$10.96 \pm 7.61$
5.	100	$60.65 \pm 1.21$	$23.23 \pm 7.19$
IC <sub>50</sub> value (µM)		82.72	> 100

Concentration of thapsigargin and ionomycin were 0.5 µM and 1 µM, respectively. Data represent mean  $\pm$  SEM of three independent experiments.

IC<sub>50</sub> of the inhibitory effects of **1** on histamine release induced by both of which were 82.72 µM and > 100 µM, respectively.

In RPMCs experiments, **1** at 100 µM induced the spontaneous histamine release by  $0.23 \pm 0.27\%$  ( $n =$

3), and this value was lower than this in RBL-2H3 cells.

## DISCUSSION

Upon tracheospasmodic guided isolation method, dihydrocubebin (**1**) was isolated from n-hexane extract of *Piper cubeba* L.f fruits traditionally used to treat respiratory disorders. It was suggested that **1** contributed anti-asthmatic activity present in the *Piper cubeba* L.f fruits. In present study, we studied the effect of dihydrocubebin on the histamine release from rat mast cell i.e. rat basophilic leukemia (RBL-2H3) cells and rat peritoneal mast cells (RPMCs).

Dinitrophenylated bovine serum albumin (DNP<sub>24</sub>-BSA) is a specific antigen for monoclonal IgE antibody (Bottcher *et al.*, 1980; Liu *et al.*, 1980). The cross-linkage of antigen into IgE antibody molecules on FcεRI receptors, and in turn can generate a series of intracellular signaling such as the activation of protein tyrosine kinases and an increase of intracellular Ca<sup>2+</sup> levels. Finally, these subsequent signaling events trigger the granule exocytosis releasing histamine from mast cells (Metcalf *et al.*, 1997). In our study, **1** markedly inhibited the histamine release from RBL-2H3 cells induced by DNP<sub>24</sub>-BSA. This finding indicates that **1** might alter the effect of DNP<sub>24</sub>-BSA on mast cells by affecting its interaction with IgE on the mast cell surface or by altering intracellular signal transduction involved in mast cell degranulation.

Furthermore, we investigated whether the inhibitory effect of 1,5-bis(4'-hydroxy-3'-methoxyphenyl)-1,4-pentadiene-3-one is related to intracellular Ca<sup>2+</sup> signal. For this purpose, we used thapsigargin and ionomycin. Thapsigargin, a sesquiterpene lactone isolated from plant *Thapsia garginica*, is able to release mediators from isolated rat mast cells (Patkar *et al.*, 1979), its target is sarco/endoplasmic reticulum Ca<sup>2+</sup> ATPase (SERCA) to prevent the Ca<sup>2+</sup> store refilling. The RE-store depletion because of intracellular Ca<sup>2+</sup> released from RE has a major

role in the opening of cell membrane Ca<sup>2+</sup> channels to cause Ca<sup>2+</sup> influx in mast cells (Metcalf *et al.*, 1997; Thastrup *et al.*, 1990). Ionomycin is a polyether antibiotic with a high affinity for calcium ions. Ionomycin is obtained in pure form from fermentation broths of *Streptomyces globatus* sp. nov. Trejo by solvent extraction (Liu *et al.*, 1978; Westley *et al.*, 1979). Ionomycin can release histamine from mast cells by increasing Ca<sup>2+</sup> intracellular from both intracellular Ca<sup>2+</sup> pools and Ca<sup>2+</sup> influx (Huang and Putney, 1998). In the study, **1** successfully inhibited the histamine release from rat mast cell induced by either thapsigargin or ionomycin. This finding indicates that the effect of this compound on the release of mast cells mediators involve mechanisms related to intracellular Ca<sup>2+</sup> signaling events or downstream processes of intracellular Ca<sup>2+</sup> signaling in mast cells.

Compound 48/80 is a histamine secretagogue which able to increase the rate of GTP S binding to G-proteins (Mousli *et al.*, 1990; Palomaki *et al.*, 2006). The activation of G-proteins can trigger intracellular signaling events such as activation of phospholipase C, protein kinase C, and Ca<sup>2+</sup> signaling, and finally results in the release of histamine from these cells. By this compound, only restricted types of mast cells are activated, and these are of the connective tissue type. Mucosal mast cell such as RBL-2H3 cells, do not respond to this secretagogue (Metcalf *et al.*, 1997). In the study, **1** did not show significant effect on compound 48/80-induced histamine release. This fact indicates that the inhibitory effect of **1** is not related to signaling events in G protein activation pathway.

Phorbol myristate acetate (PMA) stimulates histamine release by activating protein kinase C (PKC) signaling event in mast cells (Okano *et al.*, 1986; Bergstrand *et al.*, 1992). PMA does not stimulate histamine release to the same extent as other histamine secretagogue (Okano *et al.*, 1985). In order to promote granule exocytosis and mediator release from mast cells, activation of PKC is markedly enhanced in presence of a large Ca<sup>2+</sup> influx which causes

translocation of PKC to the membrane (Katakami et al., 1984; Chakravarty, 1990). In RPMCs, activation of PKC alone also caused a few amount of histamine release but the combination with Ca ionophore induced the high amount of histamine release (Heiman and Crews, 1985). A sub-effective dose of Ca<sup>2+</sup> ionophore is often used concomitantly with PMA to induce high amount of histamine release (Yen et al., 1992; Shin et al., 2004). In the study, **1** did not affect the histamine release from RMPCs induced by PMA and ionomycin in combination. It suggests that this compound did not alter the interaction between PKC and intracellular Ca<sup>2+</sup> during granule exocytotic processes.

In previous study, the n-hexane and ethanol extract of *Piper cubeba* fruit induced the histamine release in absence of antigen (Ikawati et al., 2001). In contrast, **1** that was isolated from n-hexane extract of *Piper cubeba* fruit did not induce any obvious spontaneous histamine release from either RBL-2H3 cells or RPMCs. This evidence suggests that there are other compounds in this n-hexane extract and not **1**, responsible for the spontaneous histamine release.

In conclusion, **1** inhibited histamine release and may involve in the mechanisms related to intracellular Ca<sup>2+</sup> signalings or downstream processes of intracellular Ca<sup>2+</sup> signalings in mast cells. Nevertheless, further study is required to investigate the detail mechanism of **1** in mast cells.

## REFERENCES

- Barnes PJ. (1999) Therapeutic strategies for allergic diseases. *Nature* **402**, B31-38.
- Barsumian EL, Isersky C, Petrino MG, Siraganian RP. (1981) IgE-induced histamine release from rat basophilic leukemia cell lines. *Eur. J. Immunol.* **11**, 317-323.
- Bergstrand H, Lundquist B, Karabelas K, Michelsen P. (1992) Modulation of human basophil histamine release by protein kinase C inhibitors differs with secretagogue and with inhibitor. *J. Pharmacol. Exp. Ther.* **260**, 1028-1037.
- Bottcher I, Ulrich M, Hirayama N, Ovary Z. (1980) Production of monoclonal mouse IgE antibodies with DNP specificity by hybrid cell lines. *Int. Arch. Allergy. Appl. Immunol.* **61**, 248-250.
- Brown JM, Wilson TM, Metcalfe DD. (2008) The mast cell and allergic diseases: role in pathogenesis and implications for therapy. *Clin. Exp. Allergy.* **38**, 4-18.
- Busse WW, Reed CE. (1993) Asthma : Definition and pathogenesis, In : Middleton, E., Jr., Reed CE, Ellis EF, Adkinson NF Jr., Yunginger JW, Busse WW. (Eds.). *Allergy, Principles and Practice*, vol. I, 4<sup>th</sup> Ed., Mosby, Inc., St. Louis, Missouri, p. 1173.
- Chakravarty N. (1990) The role of protein kinase C in the histamine secretion from mast cell. *Acta Physiol. Scand.* **139**, 319-333.
- Heiman AS, Crews FT. (1985) Characterization of the effects of phorbol esters on rat mast cell secretion. *J. Immunol.* **134**, 548-555.
- Hirata N, Naruto S, Inaba K, Itoh K, Tokunaga M, Inuma M, Matsuda H. (2008) Histamine release inhibitory activity of *Piper nigrum* leaf. *Biol. Pharm. Bull.* **31**, 1973-1976.
- Holt PG, Macaubas C, Stumbles PA, Sly PD. (1999) The role of allergy in the development of asthma. *Nature* **402**, B12-17.
- Huang Y, Putney JW Jr. (1998) Relationship between intracellular calcium store depletion and calcium release-activated calcium current in a mast cell line (RBL-1). *J. Biol. Chem.* **273**, 1954-1959.
- Ikawati Z, Wahyuono S, Maeyama K. (2001) Screening of several Indonesian medicinal plants for their inhibitory effect on histamine release from RBL-2H3 cells. *J. Ethnopharmacol.* **75**, 249-256.
- Katakami Y, Kaibuchi K, Sawamura M, Takai Y, Nishizuka Y. (1984) Synergistic action of protein kinase C and calcium for histamine release from rat peritoneal mast cells. *Biochem. Biophys. Res. Commun.* **121**, 573-578.
- Liu WC, Slusarchyk DS, Astle G, Trejo WH, Brown WE, Meyers E. (1978) Ionomycin, a new polyether antibiotic. *J. Antibiot. (Tokyo)* **31**, 815-819.
- Liu FT, Bohn JW, Ferry EL, Yamamoto H, Molinaro CA, Sherman LA, Klinman NR, Katz DH. (1980) Monoclonal dinitrophenyl-specific murine IgE antibody: preparation, isolation, and characterization. *J. Immunol.* **124**, 2728-2737.
- Metcalfe DD, Baram D, Mekori YA. (1997) Mast cells.

- Physiol. Rev.* **77**, 1033-1064.
- Mousli M, Bronner C, Landry Y, Bockaert J, Rouot B. (1990) Direct activation of GTP-binding regulatory proteins (G-proteins) by substance P and compound 48/80. *FEBS Lett.* **259**, 260-262.
- Nayampali SS, Satoskar RS. (1980) Evaluation of anti-allergenic activity of *Piper longum* using rat-lung perfusion technique, in Proc. 13<sup>th</sup> Annual Conf. Indian Pharmacol. Soc., Regional Res., September 30<sup>th</sup> - October 2<sup>nd</sup>, Lab. Jammu-Tawi India.
- Okano Y, Takagi H, Nakashima S, Tohmatsu T, Nozawa Y. (1985) Inhibitory action of phorbol myristate acetate on histamine secretion and polyphosphoinositide turnover induced by compound 48/80 in mast cells. *Biochem. Biophys. Res. Commun.* **132**, 110-117.
- Okano Y, Yamada K, Takagi H, Nozawa Y. (1986) Biphasic effect of phorbol myristate acetate on histamine secretion induced by compound 48/80 in rat peritoneal mast cells. *Biochem. Biophys. Res. Commun.* **137**, 1112-1118.
- Palomaki VA, Laitinen JT. (2006) The basic secretagogue compound 48/80 activates G proteins indirectly via stimulation of phospholipase D-lysophosphatidic acid receptor axis and 5-HT<sub>1A</sub> receptors in rat brain sections. *Br. J. Pharmacol.* **147**, 596-606.
- Patkar SA, Rasmussen U, Diamant B. (1979) On the mechanism of histamine release induced by thapsigargin from *Thapsia garganica* L. *Agents Actions* **9**, 53-57.
- Prabhu BR, Mulchandani NB. (1985) Lignans from *Piper cubeba*. *Phytochemistry* **24**, 329-331.
- Shin TY. (2004) *Stachys riederi* inhibits mast cell-mediated acute and chronic allergic reactions. *Immunopharmacol Immunotoxicol.* **26**, 621-630.
- Thastrup O, Cullen PJ, Drobak BK, Hanley MR, Dawson AP. (1990) Thapsigargin, a tumor promoter, discharges intracellular Ca<sup>2+</sup> stores by specific inhibition of the endoplasmic reticulum Ca<sup>2+</sup>-ATPase. *Proc. Natl. Acad. Sci. U S A* **87**, 2466-2470.
- Wahyono S. (2005) Isolation and Structure Elucidation of Tracheospasmolytic Compounds from *Piper cubeba* L.f Fruits., *Dissertation*, Gadjah Mada University, Indonesia.
- Wahyuono S, Mulyono, Wahyono, Mursyidi A. (1999) Tracheospasmolytic activity of *Piper cubeba* L.f. fructus. *Ind. J. Pharm.* **10**, 48-56.
- Westley JW, Liu CM, Evans RH, Blount JF. (1979) Conglobatin, a novel macrolide dilactone from *Streptomyces conglobatus* ATCC 31005. *J. Antibiot. (Tokyo)* **32**, 874-877.
- Wirotesangthong M, Inagaki N, Tanaka H, Thanakijcharoenpath W, Nagai H. (2008) Inhibitory effects of Piper betle on production of allergic mediators by bone marrow-derived mast cells and lung epithelial cells. *Int. Immunopharmacol.* **8**, 453-457.
- Yamatodani A, Fukuda H, Wada H, Iwaeda T, Watanabe T. (1985) High-performance liquid chromatographic determination of plasma and brain histamine without previous purification of biological samples: cation-exchange chromatography coupled with post-column derivatization fluorometry. *J. Chromatogr.* **344**, 115-123.
- Yen A, Barrett KE, Gigli I. (1992) Protoporphyrin and long-wave ultraviolet light modulate metabolic events in rat peritoneal mast cells. *J. Invest. Dermatol.* **98**, 488-493.