

## *Hypericum* source of natural antimicrobials

Pulok K Mukherjee<sup>1\*</sup>, Atul Wahile<sup>1</sup>, KFH Nazeer Ahamed<sup>1</sup> and S Rajan<sup>2</sup>

<sup>1</sup>Department of Pharmaceutical Technology, Jadavpur University, Kolkata - 700032, India; <sup>2</sup>Medicinal Plant Collection Unit, Central Council for Research in Homeopathy, Ooty 643002, TN, India

### SUMMARY

Plants of the genus *Hypericum* (Family - Hypericaceae) are herbs, shrubs or small trees and are distributed chiefly in the temperate regions of the world. About 400 different species of *Hypericum* are available throughout the globe and 20 species occur in India, including a few cultivated in gardens. Almost all plants of the genus *Hypericum* are widely used in folk medicine. Several potent phytoconstituents from different *Hypericum* species have led to the isolation of antibacterial, antifungal and cytotoxic compounds. With the development of resistance and cross resistance with different microorganisms and the evolution of so many deadly diseases the screening and evaluation of the phytoconstituents so much so the development of varied phytoconstituents for the drug development for these deadly diseases is utmost essential in every aspects. The present review on the antimicrobial use of different *Hypericum* reports the findings from and extensive literature search on the *Hypericum* species around the globe that have been assessed for antimicrobial and antiviral activity. An attempt has been made through this review to summarize the information in this aspect in order to highlight the promising species of this genus which are worthy for further investigation as leads for drug development. Over 31 different *Hypericum* species have been reported to possess such activities with their varied number of phytoconstituents. Sixteen different constituents of six different classes of phytoconstituents have been reported to be present in different varieties of *Hypericum*, which may be considered responsible for this activity.

**Key words:** *Hypericum*; Antibacterial; Antifungal; Antiviral; Phytoconstituents

### INTRODUCTION

*Hypericum* is a well known plant in herbal medicine for the therapeutic efficacy of its different species. The most potent species of this genus is *Hypericum perforatum* Linn. The genus *Hypericum* is well known for the therapeutic efficacy of its most potent species *Hypericum perforatum* Linn. Numerous compounds with documented biological activities have been reported from it (Upton, 1997).

About 400 species of *Hypericum* are spread world wide, and about 20 species are found in India (Hobbs, 1989; Anonymous, 1962). The genus *Hypericum* belongs to family Hypericaceae. These

plants are mainly small shrubs or small trees (Anonymous, 1962). Height varies from 5 cm to 3 cm. All the species shows lots of morphological variation. There was disapproval of some botanists to place genus *Hypericum* in segregate family Hypericaceae, they believe it belongs to family Guttiferae, but morphological and phytochemical differences are not sufficient to separate these two families (Robson, 1997; Taskhtajan, 1980). Almost all plants of the genus *Hypericum* are widely used in folk medicine. Several phytochemical investigations on *Hypericum perforatum* have led to the isolation of antimicrobial (Rocha *et al.*, 1995), antifungal (Khosa *et al.*, 1982) and cytotoxic compounds (Weyerstahl *et al.*, 1995). Many therapeutic properties of different components of *Hypericum perforatum* have been reported in the American Herbal Pharmacopoeia (Upton *et al.*, 1997; Bombardelli, 1995). Extracts of

\*Correspondence: Dr Pulok K Mukherjee, Department of Pharmaceutical Technology, Jadavpur University, Kolkata - 700032, India. Tel: +91-33-24181648; Fax: +91-33-2335724; E-mail: pulokm@vsnl.net

*Hypericum perforatum* have been shown to be more effective than placebo in the treatment of depression (Linde *et al.*, 1996) and different phytochemical constituents of this plant like xanthenes and flavonoid - hyperforin have been shown to be effective antimicrobials, antivirals and antibacterials against gram-positive bacteria, and to possess wound-healing potential (Bystrov, 1975; Gurevich *et al.*, 1971; Holzl *et al.*, 1989; Kitanov *et al.*, 1987; Lavie *et al.*, 1995; Bombardelli, 1995).

In India they are widely distributed in Assam, Jammu & Kashmir, Himachal Pradesh, Uttar Pradesh (Garhwal), Sikkim, West Bengal, Meghalaya, Arunachal Pradesh, Kerala and Tamilnadu (Anonymous, 1992; Asolkar *et al.*, 1992). Out of all the species available in India Nilgiri district of Tamilnadu play a major role in its content (Anonymous, 1962; Fyson, 1974). Three species are more prominent in this region viz. *Hypericum mysorensense* Wight. and Arn. (commonly known as Shrubby St. John's Wort'), *Hypericum patulum* (commonly known as 'Tumbhul' - in Bihar; 'Uralo' in Nepal; La-san-rit - in Assam; 'Paharia' - in Bengal), *Hypericum hookerianum* (commonly known as Mehandi phul in Nepal) are abundantly available every where in the Nilgiris down, near to Kotagiri and Pykara (Fyson, 1974). All of these are erect, glabrous shrub, 1-3 m height with sessile leaves, horizontal with tips curved slightly upwards, narrow elliptic lanceolate with strong midribs (Fyson, 1974; Gamble, 1984; Rajan *et al.*, 2002). Almost all the folklore claims on these three species available at Nilgiris has been scientifically proved and their psychopharmacological (Mukherjee *et al.*, 2000a), wound healing (Mukherjee *et al.*, 2000b; 2000c; 2000d), antibacterial (2001, 2002) profiles have been reported.

*Hypericum perforatum* commonly known as St. John's wort is the most popular member of the *Hypericum* family and so many reviews on its different profiles have been reported (Bombardelli, 1995). This plant was used in ancient Greece, in folklore medicines. There are many ancient superstitions regarding these herbs. Their name *Hypericum* is derived from Greek and means Over an apparition; a reference to the belief that the herb was so obnoxious to evil spirits that a whiff of it would cause them to fly. (Grieve, 1994). The naphthodianthrones,

hypericine and pseudohypericin, different flavonoids like quercetin, hyperin etc., phluroglucinols, essential oils and xanthenes has been reported to produce antidepressant, anti microbial, antioxidant, and antiinflammatory activity (Bystrov, 1975; Gurevich *et al.*, 1971; Holzl *et al.*, 1989; Kitanov *et al.*, 1987; Rocha *et al.*, 1995; Khosa *et al.*, 1982; Weyerstahl *et al.*, 1995). Different species of *Hypericum* possess antimicrobial and antiviral activities depending on their varied constituents. Based on these concepts a complete review on the various aspects of the antimicrobial usage of different species of this potent plant species has been made in this review.

### ANTIBACTERIAL POTENTIALS OF *HYPERICUM*

#### Various *Hypericum* species with antibacterial activity

The plants of *Hypericum* genus are widely used for their various significant therapeutic activities. The available plethora of literature has proved the therapeutic efficacy of these plants for their antibacterial properties which has been explained further in Table 1.

The antimicrobial activity of the leaves and stems extract of two different varieties of *Hypericum* was evaluated by the disc diffusion method. Petroleum ether, acetone, chloroform and methanol extracts of the *Hypericum mysorensense* and *Hypericum patulum* stems and leaves were investigated for their antimicrobial activity against six different strains of bacteria and fungi by zone of inhibition method (ZIM). The results showed that both the stem and leaf extracts of both species have a very broad spectrum of antibacterial activity. The petroleum ether and acetone extract of the leaves and stems of *H. mysorensense* did not show antifungal effects against any of the six fungal organisms tested but other extracts showed a potential antifungal effect which was comparable to that of griseofulvin, the standard antifungal agent. Thus these results demonstrate that the leaf and stem extracts of both the *Hypericum* species has a very broad spectrum of activity and suggest that they may be useful in the treatment of various microbial infections (Mukherjee *et al.*, 2002a). *Hypericum hookerianum* is the common ornamental plant in the garden, when chloroform, acetone and methanolic

**Table 1.** Different Hypericum species with their antimicrobial potentials

Hypericum species	Solvent used for extract	Active against	Inactive against	References
<i>H. androsaemum</i>	Ethanol-Water	SL/SA	EC	Bhakuni <i>et al.</i> (1974)
<i>H. avicularifolium</i>	Acetone	BS/EC/ST/SA	ML/CA/CU	Sakara <i>et al.</i> (1990)
	MeOH	BS	CA/CU/EC	
	CHCl <sub>3</sub>	ST/SA/ML	EC/BS	
<i>H. brasiliense</i>	Petrol	BS	-	Roch <i>et al.</i> (1995)
	CHCl <sub>3</sub>	-	BS	
<i>H. calycinum</i>	Acetone	BS/ML/ST/SA	EC/CA/CU	Sakara <i>et al.</i> (1990)
	MeOH	BS/ML/ST/SA	EC/CA/CU	
	CHCl <sub>3</sub>	BS/ML/ST/SA/EC/CU	CA	
<i>H. chinense</i>	MeOH	BS	EC	Nagai <i>et al.</i> (1987)
<i>H. cordifolium</i>	MeOH	BS/SM/SA/SF/MG/TM/V/HSV	EC/PA/ST/AF/PV	Taylor <i>et al.</i> (1995) Taylor <i>et al.</i> (1996)
<i>H. drummondii</i>	Hexane	BS/SA	-	Jayasuriya <i>et al.</i> (1983)
<i>H. elodeoides</i>	MeOH	BS/SM/SA/SF	EC/PA/HSV/PV/V/VS	Taylor <i>et al.</i> (1995) Taylor <i>et al.</i> (1996)
<i>H. typhimurium</i>	MeOH	MG/TM/AF	CA/SC	Taylor <i>et al.</i> (1995)
<i>H. empetrifolium</i>	H <sub>2</sub> O	BP-PHI, T7, T2, T4, M52	-	Delitheos <i>et al.</i> (1992)
<i>H. erectum</i>	MeOH	SA	-	Kitagaw <i>et al.</i> (1987)
<i>H. ericoides</i>	MeOH	SA	EC/KP/PA	Rios <i>et al.</i> (1987)
	CHCl <sub>3</sub>	SA/CA	EC/KP/PA	
<i>H. galioides</i>	H <sub>2</sub> O	SA/EC/TM/CA/SC	BS/PA/AN	Mechesney <i>et al.</i> (1985)
	Cyclohexane	SA/TM/AN/CA/SC	EC/BS/PA	
	Ethyl acetate	BS/SA/CA/SC	EC/PA/TM/AN	
<i>H. gentianoides</i>	H <sub>2</sub> O	BS/SA/EC/TM/CA	PA/AN/SC	McChesney <i>et al.</i> (1985)
	Cyclohexane	PA/BS/SA/SC	EC/AN/TM/CA	
	Ethyl acetate	PA/BS/SA/EC/TM/SC/CA	AN	
<i>H. hircinum</i>	H <sub>2</sub> O	SMU/SS/SA/SO/EC/PVU	PA	Barbagallo <i>et al.</i> (1987)
	MeOH	SMU/SS/SA/SO/EC	PVU/PA	
	CHCl <sub>3</sub>	PA/SMU/SS/SA/SO	EC/PVU	
	Pet. ether	SMU/SS/SA/SO/EC/PVU	PA	
<i>H. Hookerianum</i>	CHCl <sub>3</sub>	PC/BM/BC/BS/SA/EC	-	Mukherjee <i>et al.</i> (2001)
	Acetone	PC/BM/BC/BS/SA/EC	-	
	MeOH	PC/BM/BC/BS/SA/EC	-	
<i>H. japonicum</i>	ETOH	HBV/HSV	-	Minshi (1989)
<i>H. lanuginosum</i>	Acetone	BS/SA/SSO	-	Sakar <i>et al.</i> (1988)
	ETOH	BS/SA/SSO	-	
	CHCl <sub>3</sub>	SA/SSO	BS	
	Ethyl acetate	CU	-	
<i>H. maculatum</i>	H <sub>2</sub> O	CK/CT	AFL/AF/CA/CP	Chaumont <i>et al.</i> (1978)
<i>H. montbretii</i>	Acetone	SA	BS/EC/ML/ST/CA	Sakara <i>et al.</i> (1990)
	MeOH	SA	BS/EC/ML/CU/ML	
	CHCl <sub>3</sub>	BS/EC/ST/SA/CA/CU	-	
<i>H. mysorensis</i>	Pet ether	PC/BM/BC/BS/SA/EC	CA/CT/CN	Mukherjee <i>et al.</i> (2002)
	CHCl <sub>3</sub>	PC/BM/BC/BS/SA/EC/CA/CT/CN	-	
	Acetone	PC/BM/BC/BS/SA/EC	CA/CT/CN	
	MeOH	PC/BM/BC/BS/SA/EC/CA/CT/CN	-	
<i>H. origanifolium</i>	Acetone	BS/SA/SSO/CU	-	Sakara <i>et al.</i> (1988)
	CHCl <sub>3</sub>	BS/SA/SSO/EC/ST	-	
	Ethanol	BS/SSO/CU	-	
	Ethyl acetate	SA/SSO/PA	-	
<i>H. patulum</i>	MeOH	-	BS/EC	Ishii <i>et al.</i> (1984)
<i>H. patulum</i>	Pet ether	PC/BM/BC/BS/SA/EC/CA/CT/CN	-	Mukherjee <i>et al.</i> (2002)
	CHCl <sub>3</sub>	PC/BM/BC/BS/SA/EC/CA/CT/CN	-	
	Acetone	PC/BM/BC/BS/SA/EC/CA/CT/CN	-	
	MeOH	PC/BM/BC/BS/SA/EC/CA/CT/CN	-	

**Table 1.** Different *Hypericum* species with their antimicrobial potentials

Hypericum species	Solvent used for extract	Active against	Inactive against	References
<i>H. perforatum</i>	H <sub>2</sub> O	SA/SO	EC/PVU/PA/SMU	Barbagallo et al. (1987)
	Pet ether	SA/SO/PVU/PA/EC/SS	-	
	CHCl <sub>3</sub>	SA/SO	EC/PVU/PA/SS	
	MeOH	SO	EC/PVU/PA/SS/SA/SMU	
<i>H. perforatum</i>	Acetone	IV	-	Mishenkova et al. (1975); Grauds (1997); Mccutcheon et al. (1992).
	MeOH	PVU/BS/MC/SMU/MG/SA	KP/SM/AFL/AF/ TM/CA/SC/PA	
	H <sub>2</sub> O	IV/HSV/VV/SO	PV/EC/PVU/PA/SA/SMU	
	CHCl <sub>3</sub>	SA/SO	EC/PVU/PA/SS	
	Ethyl acetate	IV	-	
	Pet. ether	PVU/PA/SA/SMU/SO/SS	-	
<i>H. reflexum</i>	MeOH	SA/BS/ML/KP	PA/EC	Herrera et al. (1996)
	CHCl <sub>3</sub>	SA/ML	EC/PA	
<i>H. salsugineum</i>	CHCl <sub>3</sub>	EC/PA/SA/SSO/BS/ST/CU	-	Sakara et al. (1988)
	EtOH	EC/SSO/BS/ST/CU/SA	-	
	Ethyl acetate	SA/SSO	-	
<i>H. scabrum</i>	EtOH	SA	CA	Alshamma et al. (1979)
<i>H. triquetrifolium</i>	MeOH	BS/ST/SA/EC	ML/CA/CU	Sakara et al. (1990)
	Acetone	BS/ST/SA	EC/ML/CA/CU	
	CHCl <sub>3</sub>	BS/ST/SA/ML/CA/CU	EC	
<i>H. uliginosum</i>	EtOH-H <sub>2</sub> O	SA/TM	-	Taylor et al. (1969)

FL=Aspergillus flavus;  
 AF=Aspergillus fumigatus;  
 BM=Bacillus megaterium  
 CA=Candida albicans;  
 CP=Candida parapsilosis;  
 CK=Candida krusei;  
 EC=Escherichia coli;  
 IV=Influenza virus;  
 KP=Klebsiella pneumoniae;  
 MG=Microsporium gypseum;  
 PA=Pseudomonas aeruginosa;  
 PC=Pseudomonas cepacia  
 PV=Polio virus;  
 SL=Sarcina lutea;  
 SM=Serratia marcescens;  
 SMU=Streptococcus mutans;  
 SO=Staphylococcus oxford;  
 ST=Salmonella typhimurium;  
 VS=Virus sindbis;  
 VV=Vaccinia virus;

AN=Aspergillus niger  
 BS=Bacillus subtilis  
 BC=Bacillus coagulans  
 CN=Cryptococcus neoformans  
 CU=Candida utilis  
 CT=Candida tropicalis  
 HBV=Hepatitis B virus;  
 MC=Microsporium cookei

TM=Trichophyton mentagrophytes  
 SSO=Streptococcus sorbinus  
 SC=Saccharomyces cerevisiae  
 SA=Staphylococcus aureus  
 SF=Streptococcus faecalis  
 SS=Streptococcus sanguis  
 PVU=Proteus vulgaris  
 ML=Micrococcus luteus  
 HSV=Herpes simplex virus  
 - = No activity reported

extracts of leaves and stem were investigated against six different gram +ve and gram -ve bacteria. The stem showed maximum antibacterial activity (Mukherjee et al., 2001a). The methanolic extracts of *H. hookerianum* (Mukherjee et al., 2000b), *H. mysorens* (Mukherjee et al., 2000c) and *H. patulum* (Mukherjee et al., 2000d) have been found to possess significant wound healing activity (Mukherjee, 2002b).

The ethanolwater extract of *Hypericum androsaemum* when tested against three different bacteria it was found to have activity against *Sarcina lutea* and *S. aureus*, while inactive against *E. coli* (Bhakuni et al.,

1974). This plant also reported to have wound healing activity (Vickery, 1981). Acetone, Chloroform and Methanol extract of dried flowers and leaves of *Hypericum avicularifolium* has been reported to have activity against various bacteria like *Bacillus subtilis*, *Salmonella typhimurium*, *S. aureus* etc. while only acetone extract was active against *E. coli*. When acetone, chloroform and methanolic extracts of *Hypericum calycinum* were tested against same bacteria they were found to have activity against many of them. But only chloroform extract was found to be active against *E. coli* (Sakara et al., 1990).

Nagai *et al.* (1987) reported the activity of methanolic extract of *Hypericum chinese* flowers against *B. subtilis*, while it was found inactive against *E. coli*. The methanolic extract of fresh leaves of *Hypericum cordifolium* and methanolic extract of fresh root of *Hypericum elodeoides* were studied by Taylor *et al.* (1995a) and was reported to have activity against *B. subtilis*, *Serratia marcescens*, *S. aureus*, and *Streptococcus faecalis* while inactive against *E. coli*, *Pseudomonas aeruginosa* and *Salmonella typhimurium*. The hexane extract of dried leaf and dried root of *Hypericum drummondii* were found to be active against *Bacillus subtilis* and *S. aureus* (Jayasuria *et al.*, 1989). The methanolic extract of dried aerial parts of *Hypericum erectum* was reported to be active against *S. aureus* (Kitagawa *et al.*, 1987).

The methanolic and chloroform extracts of aerial parts of *Hypericum ericoides* when investigated for antibacterial activity, both extracts showed weak activity against *S. aureus*, while both of them was found to be inactive against *E. coli*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* (Rios *et al.*, 1987).

The cyclohexane, ethyl acetate and water extracts of dried aerial parts of *Hypericum galioides* and *Hypericum gentianoides* have been reported to have activity against various bacteria (Mc Chesney *et al.*, 1985). It was found that the activity of the extracts was mainly dependent upon solvent used for the extraction. In case of *Hypericum glioides* cyclohexane and ethyl acetate extracts were inactive against *E. coli* but aqueous extract showed activity.

Chloroform, methanol, petroleum ether and aqueous extracts of dried aerial parts of *Hypericum hircinum* has been evaluated for their antibacterial potentials and most of these extracts showed activity against *Streptococcus mutants*, *S. sanguis*, *S. aureus*, *S. oxford* and *E. coli*. When same solvent extracts of dried aerial parts of *Hypericum perforatum* were evaluated, only petroleum ether extract showed activity against maximum number of strains of bacteria (Barbagallo *et al.*, 1987). *Hypericum lanuginosum*, *Hypericum origanifolium* and *Hypericum salsugineum* showed activity against *S. aureus*, *S. sorbinus* and *B. subtilis*, when extracted with acetone, chloroform, ethyl acetate and ethanol (Sakara *et al.*, 1988). *Hypericum montbret* and *Hypericum triquetrifolium* showed antibacterial activity against *S. aureus*, *B.*

*subtilis*, *E. Coli* and *S. typhimurium* when dried leaves and flowers were extracted with chloroform, acetone and methanolic extracts of some plant showed activity against only *S. aureus* (Sakara *et al.*, 1990). Though the methanolic extract of *Hypericum patulum* is inactive against *B. subtilis* and *E. coli*, (Ishii *et al.*, 1984), it showed wound healing activity in excision and incision models in rat (Mukherjee *et al.*, 2000d).

Methanolic and chloroform extracts of entire plant *Hypericum reflexum* showed activity against *S. aureus*, *S. epidermidis*, *B. cereus*, *B. subtilis* and *Micrococcus luteus* (Herrera *et al.*, 1996). Ethanol extract of dried aerial parts of *Hypericum scabrum* showed antibacterial activity against *S. aureus* (Al-shamma *et al.*, 1979). *Hypericum uliginosum* showed antibacterial activity against *S. aureus*, when extracted with ethanol water (Taylor *et al.*, 1969).

*Hypericum perforatum* is the most popular species of the genus beside its significant antidepressant and other pharmacological actions, its antimicrobial activities are also reported. Its various extracts showed antibacterial action against different strains of microorganisms, these activities are summarized in Table 1.

#### **Different phytoconstituents of Hypericum species with antibacterial potential**

Numerous compounds with documented biological activities has been reported from *H. perforatum* e.g. naphthodianthrones, hypericin and pseudohypericin, different flavonoids like quercetin, hyperin etc., phloroglucinols, essential oils and xanthenes has been reported to produce antidepressant, anti microbial, antioxidant, and anti-inflammatory activity (Upton, 1997; Bystrov, 1975; Gurevich *et al.*, 1971; Holzl *et al.*, 1989; Kitanov *et al.*, 1987; Rocha *et al.*, 1995; Khosa *et al.*, 1982; Weyerstahl *et al.*, 1995). The different extracts as well as the different phytochemical constituents of this plant like xanthenes and the flavonoid - hyperforin have been shown to be effective as antiviral and antibacterial against gram positive bacteria and possess wound healing potentials (Lavie *et al.*, 1995; Bombardelli, 1995).

Thus various species of *Hypericum* have been reported to possess antibacterial activity and many of them were reported to have potentials antibacterial activities comparatively at lower concentrations.


**Table 2.** Phytoconstituents responsible for antimicrobial and antiviral activity from various species of *Hypericum*

Constituents	Activity	Plants	Reference
Phloroglucinol	Antibacterial	<i>H. perforatum</i>	Bystrov 1975; Gurvich et al., 1971
		<i>H. japonicum</i>	Yamaki et al., 1994; Ishiguro et al., 1994
		<i>H. brasiliense</i>	Rocha et al., 1995, 1996
		<i>H. hirsutum</i>	Umek et al., 1990
		<i>H. maculatum</i>	
		<i>H. tetrapetum</i>	
		<i>H. montanum</i>	
		<i>H. humifusum</i>	
Xanthonenes	Antibacterial, antifungal and Antiviral	<i>H. perforatum</i>	Upton, 1997
		<i>H. brasiliense</i>	Rocha et al., 1994
		<i>H. patulum</i>	Ishiguro et al., 1993
		<i>H. patulum</i>	Ishiguro et al., 1995a,b
		<i>H. androsaemum</i>	Niesen et al., 1979
Hypericine	Antiviral	<i>H. perforatum</i>	Lavie et al., 1995; Meruelo et al., 1988
		<i>H. hirsutum</i>	Umek et al., 1994;
		<i>H. maculatum</i>	Karting et al., 1996;
		<i>H. tetrapetum</i>	Brantner et al., 1994
		<i>H. montanum</i>	
		<i>H. humifusum</i>	
		<i>H. tomentosum</i>	
		<i>H. bithynicum</i>	
		<i>H. glandulosum</i>	
Pseudo hypericin	Antiviral	<i>H. perforatum</i>	Meruelo et al., 1988
		<i>H. hirsutum</i>	Umek et al., 1994;
		<i>H. maculatum</i>	Karting et al., 1996;
		<i>H. tetrapetum</i>	Brantner et al., 1994
		<i>H. montanum</i>	
		<i>H. humifusum</i>	
		<i>H. tomentosum</i>	
		<i>H. bithynicum</i>	
		<i>H. glandulosum</i>	
Proanthocyanidin	Antibacterial, antifungal and antiviral	<i>H. perforatum</i>	Upton 1997
Essential oils	Antifungal	<i>H. perforatum</i>	Khosa et al., 1982

Various phytoconstituents of *H. perforatum* responsible for antimicrobial potentials has been shown in Table 2 and few phytoconstituents has been shown in Fig. 1. Hyperforin (I) a phloroglucinol derivative has potent antibacterial activity against gram positive bacteria and also reported to have wound healing activity (Bystrov, 1975; Gurevich et al., 1971). Xanthonenes are also reported to have antimicrobial activity. Yamaichi et al. (1994) reported the antimicrobial activity of Phloroglucinol isolated

from *Hypericum japonica*, against *Staphylococcus aureus*. Ishiguro et al. (1987) reported saroapidin A, B, and C, (II) isolated from *Hypericum japonicum* as antibiotic compounds, same author in 1990 reported the antimicrobial activity of sarothrali G, isolated from *Hypericum japonicum*.

Ishiguro and co-workers (1994) isolated two new phloroglucinol derivatives viz. sarothralen C and D (III) from the methanolic and ether extract of whole plant of *Hypericum japonicum*, which has



**Fig. 1.** Phytoconstituents of *Hypericum* Species with antimicrobial potentials.

been reported to have antibacterial activity against *S. aureus*. Four phloroglucinol derivatives viz. Japonicine-A (IV), uliginosin-A, isouliginosin-B (V) and hyperbrasilol-A, isolated from *Hypericum brasiliense* petrol extract are reported to have activity against *Bacillus subtilis*. New phloroglucinol viz. hyperbrasilol B has been reported from the petrol extract of leaves and flowers of *Hypericum*

*brasiliense*, and three new phloroglucinol were found to have antibacterial activity against *B. subtilis* (Rocha *et al.*, 1995; 1996).

The xanthenes compounds are responsible for the antimicrobial activities of *Hypericum perforatum* (Upton, 1997). The literature survey shows the presence of xanthenes in many other species of *Hypericum* viz. *Hypericum paturum* (Ishiguro *et al.*,

1993), *Hypericum patulam* (Ishiguro et al., 1995a; 1995b), *Hypericum androsaemum* (Nielsen et al., 1979). Hyperforin which is a well proved antibacterial phytoconstituents was also found to be present in many *Hypericum* species like *Hypericum perforatum*, *Hypericum hirsutum*, *Hypericum maculatum*, *Hypericum tetrapetum*, *Hypericum montanum*, and *Hypericum humifusum* (Umek et al., 1990) which are responsible for the antibacterial activity of other *Hypericum* species except *H. perforatum*.

### ANTIFUNGAL POTENTIALS OF *HYPERICUM*

#### Different *Hypericum* species with antifungal activity

Various plant extracts of *Hypericum* has been reported to have antifungal potentials. The potential species for this purpose has been explained in Table 1. The acetone, chloroform and methanol extracts of dried flowers and leaves of *Hypericum avicularifolium*, *Hypericum calycinum*, *Hypericum montbretii* and *Hypericum triquetrifolium* when investigated for anti-yeast activity, only chloroform extracts were found to be active against *Candida utilis* (Sakara et al., 1990). Methanolic extract of fresh leaves of *H. cordifolium* showed antifungal activity against *Microsporum gypseum* and *Trichophyton mentagrophytes*. The activity was enhanced with exposure to UV light (Taylor et al., 1995a). *Hypericum gentianoides* and *Hypericum galioides* showed antifungal activity against various fungi strains, when extracted with cyclohexane and ethyl acetate (Mc Chesney et al., 1985). Aqueous extract of dried aerial parts of *Hypericum hirsutum* was proved active against *Candida krusei*, *Candida parapsilosis*, *C. pseudotropicalis* and *C. tropicalis*, while aqueous extract of *H. maculatum* has activity against *C. Krusei* and *C. tropicalis* (Chaumont et al., 1978a).

Antiyeast activity of *H. lanuginasum*, *H. organifolium* and *H. salsugineum* has been reported by Sakara et al. (1988). Aqueous extract of fresh entire plant of *H. pilchard* is reported to have antifungal activity against pathogenic fungi *Pestalotia funerea*, *Aclerotinia trifoliorum* and *Stereum purpureum* (Chaumont et al., 1978b). *Hypericum revolutum* is reported as active against various fungi (Decosterd et al., 1987).

Methanolic extract of *Hypericum perforatum* has

been reported to possess weak activity against *Microsporum cookei* and *Microsporum gypsum* but ethanol extract has been reported activity against various other fungi. (Mc Cutecheon et al., 1994; Khosa et al., 1982).

#### Different phytoconstituents of *Hypericum* species with antifungal potential

The antifungal activities of *Hypericum perforatum* are supposed to be due to the essential oil. The essential oil of this plant contain monoterpenes and sesquiterpene (caryophyllen, humulene) (Khosa et al., 1982; Sticher, 1977; Chialva, 1981; Mathis, et al., 1964). The oil is also said to have the use as sunscreen, which is disputed. The oil has been used in commercial sunscreen products. But reports of its efficacy are contradictory (Proserpio, 1976; Anonymous, 1992). Hyperbrasilone (VI) which is  $\gamma$ -pyrone and various xanthenes isolated from the *Hypericum brasiliense* has been shown anti-fungal activity against *Cladosporium cucumerrinum* (Rocha et al., 1994). The antifungal properties of different phytoconstituents of *Hypericum* have been explained in Table 2 and few phytoconstituents has been shown in Fig. 1. The various xanthone derivatives have been also reported to have antifungal potentials (Upton, 1997). More investigations are required to find out the antifungal activity of various phytoconstituents occurs in different species of this genus, against various pathogenic fungi.

### ANTI VIRAL SPECTRUM OF DIFFERENT SPECIES OF *HYPERICUM*

#### Different *Hypericum* species with antiviral potential

St. John's Wort' or *H. perforatum* is the popular member of genus *Hypericum* which is being used from the time of the ancient Greeks for its varieties of Pharmacological and antimicrobial activities. Many writers like Hippocrates, Pliny, Dioscorides, Theophrastus and Galen has mentioned about its medicinal properties (Upton, 1997). It is very popular as a antidepressant in America and European countries. This plant has got more importance due to the presence of hypericin, which is said to have antiviral activity against HIV infestation.

*Hypericum perforatum* showed antiviral activity against *Influenza virus* (Mishenkova, 1975), *Herpes*

*simplex* virus type 2 and *Vaccinia virus* (May *et al.*, 1978). When hypericin isolated from *H. perforatum* was investigated as a chemotherapeutic agent for HIV infection, it was found effective in inhibition of replication cycle of virus. (Vilietinck *et al.*, 1998). When 18 patients with acquired immunodeficiency syndrome (AIDS) were treated with intra-venous *Hypericum* preparation and tablets of different dosage 16 patients out of 18 responded the treatment and 14 patients out of those 16 remained clinically stable (Steinbeck *et al.*, 1993). Though hypericin has given the hope for treatment of HIV infection, the proper concentration required for efficacy is not known. In large dosage it may cause photosensitization as has been reported in case of various animals, which are either grazing on *Hypericum* or given the dosage of hypericin (Munscher, 1961; Southwell *et al.*, 1991; Horsley, 1934).

From past many years the antiviral activities of various species of this genus has been investigated and many of them are proved to have antiviral activities. The dried roots juice *H. cardifolium* and *H. uralum* has shown activity against *Polio virus -1*, *Herpes simplex-1* and *Sindbis virus* (Taylor *et al.*, 1996). Aqueous and ethanol extracts of *Hypericum japonicum* showed strong activity against *Hepatitis B virus* and *Herpes simplex 2 virus* (Minshi 1989, Zheng *et al.*, 1992).

#### **Different phytoconstituents of *Hypericum* species with antiviral potential**

Many authors have reported the antiviral activities of hypericin (VII) and pseudohypericin (VIII), which are the phytoconstituents present in the *H. perforatum* and many other species of *Hypericum* (Meruelo *et al.*, 1988; Weber *et al.*, 1994; LopezBazzocchi *et al.*, 1991; Wood *et al.*, 1990). Xanthonenes present in the *Hypericum perforatum* also have antiviral potentials (Upton, 1997; Hobbs, 1989) various other xanthonenes are present in many other species of *Hypericum*, so it is needed to investigate their possible antiviral potentials of those phytoconstituents like xanthonenes, hypericin and pseudohypericin etc. which are reported to be present in many other varieties viz. *H. hirsutum*, *H. maculatum*, *H. tetrapterum*, *H. montanum*, *H. humifusum*, *H. tomentosum*, *H. bithynicum*, *H. glandulosum* and *H. balearicum* (Umek *et al.*, 1994). The antiviral potentials of various

*Hypericum* species may be due to presence of hypericin, pseudohypericin and xanthonenes, to find out the possible anti-viral potentials of various species of *Hypericum*, more phytochemical investigations and antiviral activity studies are required. The antiviral potentials of various phytoconstituents of different *Hypericum* species have been shown in Table 2 and few phytoconstituents has been shown in Fig. 1.

## **CONCLUSIONS**

The resurgence of the microbial and viral diseases has been known to occur in its various resistant from time to time in endemic and epidemic proportions all over the world. From time immemorial the cure of these diseases has been attempt by the traditional medicine. Scientific validation and screening for new phytotherapeutic agents from this traditional knowledge has helped for the drug development with the emergence of resistance and cross resistance to some standard drugs used in these particular infections. Over 31 different *Hypericum* species and various compounds derived from them have been evaluated for their antibacterial, antifungal and anti viral potentials. Composite or non-specific activities of plant extracts are apparently significant in the uses and cure with traditional knowledge. The data compiled in Table 1 reveal that extracts of different *Hypericum* species from the same family have shown varied anti microbial and antiviral potentials. Table 2 showed the potent plant derived phytoconstituents from this particular species responsible for these activities. This review on the combined approaches of exploration and exploitation of the different species of *Hypericum* for their uses in the field of bacterial and viral infection will definitely throw further light to different findings for the exploitation of the lead compounds for treating the deadly ailments and in one hand and to restore confidence in the use of *Hypericum* in the other.

## **ACKNOWLEDGEMENTS**

The authors are thankful to The All India Council for Technical Education (AICTE), Govt. of India, New Delhi for financial support through AICTE.

Dr. PKM is thankful to Librarian of the Bibliothek, Leiden/Amsterdam Center for Drug research for providing the complete searching facilities. Thanks are due to the Department of Science & Technology, Govt. of India for financial support.

## REFERENCES

- Alshamma A, Mitscher LA. (1979) Screening results of 327 species for alkaloids and antimicrobial agents. *J. Natural Product* **42**, 633-642.
- Anonymous. (1962) *The Wealth of India*, Vol. V, p. 154, Publication & Information Directorate CSIR, New Delhi, India.
- Anonymous. (1992) *Dictionary of Indian medicinal plants*, P. 252, CIMAP, Lucknow.
- Asolkar LV, Kakkar KK and Chakre OJ. (1992) *Second Supplement to Glossary of Indian Medicinal Plant with Active Principles*, Vol. 1, pp. 138-139 Publication and Information Directorate, CSIR, New Delhi.
- Barbagallo C, Chisari G. (1987) Antimicrobial activity of three *Hypericum* species. *Fitoterapia* **58**, 175-177.
- Bhakuni DS, Bittner M, Marticorenac, Silva M, Weldt E, Melo ME. (1974) Screening of Chilean plants for antimicrobial activity. *L. Loydia* **37**, 621-632.
- Bombardelli E, Morazzoni P. (1995) *Hypericum perforatum*. *Fitoterapia* **66**, 43-68.
- Bronz I. (1983) The absolute configuration of hyperforin, an antibiotic from *H. perforatum* L; based on the crystal structure determination of its p- bromobenzoate ester. *Acta. Chem. Scand.* **37**, 263-265.
- Bystrov NS, Chernov BK and Dobrynin VN. (1975) The structure of hyperforin. *Tetrahedron Lett.* **32**, 2791-2794.
- Branter A, Kartring TH, Quehenberger F. (1994) Vergleichende phytochemische untersuchungen an *H. perforatum* and *H. maculatum* crantz. *Scientia Pharmaceutica* **62**, 261-276.
- Chaumont JP, Senet JM. (1978a) Antagonistic properties of higher plants fungal parasites of man from food contaminants screening of 200 fungi. *Planta Med.* **12**, 186-196.
- Chaumont JP, Bourgeois M. (1978b) Antagonistic properties of one hundred plant extracts on 7 pathogenic fungi. *L. Loydia* **41**, 437-441.
- Chialva F. (1981) Study on the composition of the essential oil from *H. perforatum* L. and *Teucrium chamaedrys* L. Review. *EPPOS* **63**, 286-288.
- Delitheos A, Papadimitriou C, Yannitsaros A. (1992) Investigation of antiphage activity in plant extracts. *Fitoterapia* **63**, 441-450.
- Decosterd LA, Stoeckli E, Msoutthi JD. (1987) New antifungal chromenyl ketones and their pentacyclic dimers from *H. revolutum*. *Helvetica Chim. Acta.* **70**, 1694-1702.
- Fyson PF. (1974) *The Flora of the Nilgiris and Pulney Hill Tops*, Vol. 1, pp. 37-39, Bishen Singh & Mahendra Pal Singh, Dehradun.
- Grauds C. (1997) St. Johns wort for depression. *Pharmacy Times* **63**, 40.
- Gamble JS. (1984) *Flora of the Presidency of Madras*, Vol. 1, pp. 69-71, Bishen Singh & Mahendra Pal Singh, Dehradun.
- Grieve M. (1994) *A Modern Herbal*, p. 708, Tiger books international, London.
- Gurevich AI. (1971) Hyperforin, an antibiotic from *H. perforatum*. *Antibiotiki* **16**, 510-513.
- Hobbs C. (1989) St. Johns wort: *H. perforatum* L. A Review. *Herbal Gram* **18/19**, 24-33.
- Herrera RM, Perez M, Merrera DA. (1996) Antimicrobial activity of extracts from plants endemic to the canary islands. *Phytother. Res.* **10**, 364-366.
- Holz J, Demisch L, Gollnik B. (1989) Investigation about anti-depressive and mood changing effects of *Hypericum perforatum*. *Planta Med.* **55**, 643.
- Horsley CH. (1934) Investigation in to the action of St. Johns wort. *J. Pharmacol. Exp. Ther.* **50**, 310.
- Ishiguro K, Yamaki M, Kashihara M. (1987) Saroaspidin A, B & C, additional antibiotic compounds from *H. japonicum*. *Planta Med.* **53**, 415-417.
- Ishiguro K, Nagata S, Fukumoto H, Yamaki M, Isoi K. (1994) Phloroglucinol derivatives from *H. japonicum*. *Phytochemistry* **35**, 469-471.
- Ishiguro K, Nakajima M, Fukumoto H, Isoi K. (1995b) Xanthone substituted with anirregular monoterpene in cell suspension culture of *Hypericum patulum*. *Phytochemistry* **39**, 903-905.
- Ishiguro K, Yamaki M, Kashihara M, Takagi S, Isoi K. (1990) Sarothralin G, a new antimicrobial compound from *H. japonicum*. *Planta Med.* **56**, 274-276.
- Ishiguro K, Fukumoto H, Nakajima M, Isoi K. (1993) Xanthenes in cell suspension cultures of *H. patulum*. *Phytochemistry* **33**, 839-840.
- Ishiguro K, Nagata S, Fukumoto H, Yamaki M, Isoi K. (1994) Pluroglucinol derivatives from *H. japonicum*. *Phytochemistry* **35**, 469-471.
- Ishiguro K, Nakajima M, Fukumoto H, Isoi K. (1995a) Co-occurrence of prenylated xanthenes and their cyclization products in cell suspension cultures of *H. patulum*. *Phytochemistry* **38**, 867-869.
- Ishii R, Yoshikawa K, Minakata H, Komura H, Kada T. (1984) Specificity of bio-antimutagens in plant kingdom. *Agricultural Biological Chemistry* **48**, 2587-

- 2591.
- Jayasuria H, Mechesney JD, Swanson SM, Pezzuto JM. (1989) Antimicrobial and cytotoxic activity of rottlerin type compounds from *H. drummonds*. *Journal of Natural Product* **52**, 325-331.
- Kartnig T, Gobel I, Heydel B. (1996) Production of Hypericin, Pseudohypericin and Flavonoids in cell cultures of various *Hypericum* species. *Planta Med.* **62**, 51-53.
- Kitagawa S, Nishibe S, Baba H. (1987) Studies of the Chinese crude drug. *Yakugakuzasshi* **107**, 274-278.
- Khosa RL, Bhatia N. (1982) Antifungal effect of *H. perforatum* L. *Journal Scientific Research Plant Medicine* **3**, 49-50.
- Kitanov GM, Blinova KF. (1987) Modern state of the chemical study of species of the genus *Hypericum*. *Khimiya Prirodnykh Soedinenni* **2**, 185-203.
- Lavie J, Mazur Y, Lavie D and Meruelo D. (1995) The chemical and biological properties of hypericin- a compound with a broad spectrum of biological activities. *Medicinal Research Review* **15**, 111-119.
- Linde K, Gilbert R, Mulrow C D, Pauls A, Weidenhammer W, Melchart M. (1996) St. Johns Wort for depression an overview and meta-analysis of randomised clinical trials. *Brit. Med. J.* **313**, 253-257.
- Lopez-Bazzocchi I, Hudson J B, Towers GHN. (1991) Antiviral activity of the photo active plant pigment Hypericin. *Photochemical Photobiology* **54**, 95-98.
- Mchesney JD, Adams RP. (1985) Co-evaluation of plant extracts as petrochemical substitutes and for biologically active compounds. *Econ. Bot.* **39**, 74-86.
- Meruelo D, Lavie G, Lavie D. (1988) Therapeutic agents with dramatic antiretroviral activity and little toxicity at effective doses: aromatic polycyclic diones hypericin and pseudohypericin. *Proceedings National Academy of Science* **85**, 5230-5234.
- Minshi Z. (1989) An experimental study of the anti-HSV II action of 500 herbal drugs. *Journal of Traditional Chinese Medicine* **9**, 113-116.
- Mishenkova EL, Derbentseva NA, Garagulya AD. (1975) Antiviral properties of St. Johns Wort and preparations produced from it. *Tr. sezda. Miterobiol. Ukr.* **4**, 222.
- Mathis C and Ourisson G. (1964) Etude chimio taxonomique du genre *Hypericum* IV. *Phytochemistry* **3**, 377-378.
- May G, Willuhn G. (1978) Antiviral activity of aqueous extracts from medicinal plants in tissue culture. *Arznein-forsch* **28**, 1-7.
- McCutecheon AR, Ellis SM, Hancock REW, Towers GHN. (1994) Antifungal screening of medicinal plants of British columbian native peoples. *J. Ethnopharmacol.* **44**, 157-169.
- Munscher WC. (1961) *Poisonous plants of the United states*, The Macmillan co., New York.
- Minshi Z. (1989) An experimental study of the anti-HSV II action 500 herbal drugs. *Journal of Traditional Chines Medicine* **9**, 113-116.
- Mishenkova EL, Derbentseva NA, Garagulya AD, Litvin LN. (1975) Antiviral properties of St. Johns wort and preparations produced from it. *Tr. Sezda.mikrobiol. Ukr.* **4**, 222.
- Mukherjee PK, Suresh B, Verpoorte R. (2000a) CNS active potentials of some *Hypericum* species of India. *Phytomedicine* **8**, 331-337.
- Mukherjee PK, Suresh B. (2000b) The evaluation of wound healing potential of *H. hookerianum* leaf and stem extracts. *J. Altern. Complem. Med.* **6**, 61-69.
- Mukherjee PK, Suresh B. (2000c) Studies on in vivo wound healing activity of leaf extract of *H. mysorensis* with different wound models in rat. *Natural Product Sciences* **6**, 73-78.
- Mukherjee PK, Verpoorte R, Suresh B. (2000d) Evaluation of in vivo wound healing activity of *H. patulum* leaf extract on different wound models in rats. *J. Ethnopharmacol.* **70**, 315-321.
- Mukherjee PK, Saritha GS, Suresh B. (2001) Antimicrobial spectrum of *Hypericum hookerianum*. *Fitoterapia* **72**, 558-560.
- Mukherjee PK, Saritha GS, Suresh B. (2002a) Antimicrobial potentials of two different *Hypericum* species available in India. *Phytother. Res.* **16**, 692-695.
- Mukherjee PK (2002b). *Quality Control of Herbal Drugs*, pp. 549-592, Business Horizons, New Delhi, India.
- Nagai M, Tada M. (1987) Antimicrobial compounds, chinesisin I and II from flowers of *H. Chinese*. *Chem. Lett.* **7**, 1337-1340.
- Nielsen H, Arends P. (1979) Xanthones constituents of *H. androsaemum*. *Journal of Natural Products* **42**, 301-304.
- Proserpio G. (1976) Natural sunscreens vegetable derivatives as sunscreens and tanning agents. *Cosmetics Toiletries* **91**, 34-46.
- Rajan S, Sethuraman M, Mukherjee PK. (2002) Ethnobiology of Nilgiri Hills, India. *Phytother. Res.* **16**, 98-116.
- Rios JS, Recio MC, Villar A. (1987) Antimicrobial activity of selected plants employed in the spanish mediterranean area. *J. Ethnopharmacol.* **21**, 139-152.
- Robson NKB. (1997) Bulletin British (Natural history). *Botany* **5**, 293.
- Rocha L, Marston A, Kaplan AC, Evans HS, Thull U.

- (1994) An antifungal gamma pyrone and xanthenes with MAO inhibitory activity from *H. brasiliense*. *Phytochemistry* **36**, 1381-1385.
- Rocha L, Marston A, Potterat O, Auxiliadora M, Kaplan C, Evans HS, Hostettmann K. (1995) Antibacterial phloroglucinols and flavonoids from *H. Brasiliense*. *Phytochemistry* **40**, 1447-1452.
- Rocha L, Marston A, Potterat O, Kaplan M, Hostettmann K. (1996) More phloroglucinols from *Hypericum brasiliense*. *Phytochemistry* **42**, 185-188.
- Sakar MK, Tamer AU, Tokur S. (1988) Antimicrobial activities of some *Hypericum* species growing in Turkey. *Fitoterapia* **59**, 49-52.
- Sakara MK, Tamer AU. (1990) Antimicrobial activity of different extracts from some *Hypericum* species. *Fitoterapia* **61**, 464-466.
- Southwell IA, Campbell MH. (1991) Hypericin content variation in *H. perforatum* in Australia. *Phytochemistry* **30**, 475-478.
- Steinbeck KA, Wernet P. (1993) Successful long term treatment over 40 months of HIV patients with intravenous *Hypericum*, International Conference on AIDS, Germany.
- Sticher O. (1977) Plant mono-, di-, and sesquiterpenoides with pharmacological or therapeutic activity. In: *New natural products and plant drugs with pharmacological, biological or therapeutical activity*, Edited by Wagner H and Wolff P, Springer-Verlag publication, New York.
- Taskhtajan AL. (1980) *Botany Review* **46**, 225.
- Taylor RS, Manadhar NP, Towers GHN. (1995a) Screening of selected medicinal plants of Nepal for antimicrobial activities. *J. Ethnopharmacol.* **46**, 153-159.
- Taylor RS, Mandhar NP, Hunson JB, Towers GHN. (1995b) Antiviral activities of Nepalese medicinal plants. *J. Ethnopharmacol.* **46**, 133-152.
- Taylor RS, Mandhar NP, Hudson JB, Towers GHN. (1996) Antiviral properties of Nepalese medicinal plants. *J. Ethnopharmacol.* **52**, 157-163.
- Upton R. (1997) *American Herbal Pharmacopoeia and Therapeutic Compendium, St. Johns Wort*, p. 12, Pittsburgh, PA, USA.
- Umek A, Kreft S, Kartnig T, Heydel B. (1999) Quantitative phytochemical analysis of six *Hypericum* species growing in Slovenia. *Planta Med.* **65**, 388-390.
- Vickery AR. (1981) Traditional uses and folklore of *Hypericum* in the British isles. *Economic Botany* **35**, 289-295.
- Vleitnick AJ, DeBruyne T, Apers S, Pieters LA. (1998) Plant derived leading compounds for the chemotherapy of HIV infection. *Planta Med.* **64**, 92-109.
- Taylor HL, Brooker RM. (1969) The isolation of Uliginodin A and B from *H. uliginosum* L. *Lloydia* **32**, 217-219.
- Weyerstahh P, Splittegerber U and Marschall H. (1995) Constituents of the leaf essential oil of *Hypericum perforatum* L. from India. *Flavours and Fragrance Journal* **10**, 365-370.
- Wood S, Huffman J, Weber N, Andersen D, North J, Murray B, Sidwell R, Hughes B. (1990) Antiviral activity of naturally occurring anthraquinones and anthraquinone derivatives. *Planta Med.* **56**, 651-652.
- Weber ND, Murray BK, North JA, Wood SG. (1994) The antiviral agent hypericin has *in-vitro* activity against HSV-1, through non specific association with viral and cellular membranes. *Anti viral chemistry and Chemotherapy* **5**, 83-90.
- Yamaki M, Miwa M, Ishiguro K, Takagi S. (1994) Antimicrobial activity of naturally occurring and synthetic phloroglucinol against *Staphylococcus aureus*. *Phytother. Res.* **8**, 112-114.
- Zheng MS, Zheng YF. (1992) Experimental studies on inhibition effects of 1000 Chinese medicinal herbs on the surface antigen of Hepatitis B virus. *Chung ITSA China* **12**, 193-195.